

10-1-2004

Biofilm Formation in Plant-Microbe Associations

Bronwyn E. Ramey

Indiana University, Bloomington IN

Maria Koutsoudis

University of Connecticut, Storrs CT

Susanne B. von Bodman

University of Connecticut, susanne.vonbodman@uconn.edu

Clay Fuqua

Indiana University, Bloomington IN

Follow this and additional works at: http://digitalcommons.uconn.edu/plsc_articles

 Part of the [Plant Sciences Commons](#)

Recommended Citation

Ramey, Bronwyn E.; Koutsoudis, Maria; von Bodman, Susanne B.; and Fuqua, Clay, "Biofilm Formation in Plant-Microbe Associations" (2004). *Plant Science Articles*. 18.

http://digitalcommons.uconn.edu/plsc_articles/18



ELSEVIER

Biofilm formation in plant–microbe associations

Bronwyn E Ramey¹, Maria Koutsoudis², Susanne B von Bodman^{2,3} and Clay Fuqua^{1*}

Bacteria adhere to environmental surfaces in multicellular assemblies described as biofilms. Plant-associated bacteria interact with host tissue surfaces during pathogenesis and symbiosis, and in commensal relationships. Observations of bacteria associated with plants increasingly reveal biofilm-type structures that vary from small clusters of cells to extensive biofilms. The surface properties of the plant tissue, nutrient and water availability, and the proclivities of the colonizing bacteria strongly influence the resulting biofilm structure. Recent studies highlight the importance of these structures in initiating and maintaining contact with the host by examining the extent to which biofilm formation is an intrinsic component of plant–microbe interactions.

Addresses

¹ Department of Biology, 1001 East 3rd Street, Jordan Hall 142, Indiana University, Bloomington, Indiana 47405, USA

² Molecular and Cell Biology Department, and ³ Plant Science Department U163, 1390 Storrs Road, ABL 302B, University of Connecticut, Storrs, CT 06269-4163, USA

*e-mail: cfuqua@bio.indiana.edu

Current Opinion in Microbiology 2004, 7:602–609

This review comes from a themed issue on Growth and development
Edited by Mike Tyers and Mark Buttner

Available online 27th October 2004

1369-5274/\$ – see front matter

© 2004 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.mib.2004.10.014

Abbreviations

DSF diffusible signal factor

EPS extracellular polymeric substances

GFP green fluorescent protein

LapA large adhesion protein A

QS quorum-sensing

Xcc *Xanthomonas campestris* pv. *Campestris*

Xf *Xylella fastidiosa*

Introduction

Many microorganisms in the natural environment exist in multicellular aggregates generally described as biofilms, associated with solid surfaces and in intimate contact with other microbial cells [1–3]. Cells adhere to surfaces and each other through a complex matrix comprising a variety of extracellular polymeric substances (EPS) including exopolysaccharides, proteins and DNA. Biofilm configurations range in complexity from flat, relatively


featureless films, to tightly clustered aggregates, to complex heterogeneous cellular arrangements such as towers and streamers. Cells within biofilms are physiologically distinct from the same cells grown in dispersed culture [4,5]. Biofilm cells respond to nutrient and waste product diffusion gradients, modulate their metabolism as a function of their position within the biofilm, contact adjacent cells, and engage in cell–cell communication. Adherent populations exhibit elevated antimicrobial tolerance as a consequence of biofilm structure and physiological adaptation [3]. Biofilms have tremendous practical importance in industrial, medical and agricultural settings, exhibiting both beneficial and detrimental activities.

Although most fundamental work on microbial biofilms has focused on abiotic surfaces, it is clear that biofilms can and do form on biotic surfaces during host–microbe interactions [2]. Most plant–bacterial associations rely upon the physical interaction between bacteria and plant tissues. Direct observations of bacteria adhered to plant surfaces have revealed multicellular assemblies variably described as microcolonies, aggregates and cell clusters [6,7,8]. These multicellular structures exhibit many of the defining attributes of biofilms — groups of cells enmeshed within an EPS matrix on a solid surface. For the purposes of this review, we will therefore refer to these different multicellular structures as biofilms, bearing in mind their differences from the paradigmatic microbial biofilm, and highlighting their unique properties. Our focus will be on recent work regarding the structure, formation and activity of microbial biofilms associated with terrestrial plants.

Plant surfaces: complex and dynamic environments

The terrestrial environment harbors abundant and diverse microbial populations that can compete for and modify resource pools. In this complex and competitive environment, plants offer protective oases of nutrient-rich tissues. Plants are colonized by bacteria on their leaves, roots, seeds and internal vasculature (Figure 1). Each tissue type has unique chemical and physical properties that represent challenges and opportunities for microbial colonists. Biofilms may form upon association or at later stages, with significant potential to direct or modulate the plant–microbe interaction. Additional temporal and spatial complexity arises as many microbes actively modify the colonized plant environment.

Figure 1



Sites of microbial colonization on terrestrial plants. Stylized dicotyledonous plant are depicted. Phyllosphere/phyloplane on leaves, rhizosphere/rhizoplane on roots, and internal vascular system are highlighted. Figure adapted and reproduced with permission from William C Brown Publishers (McGraw-Hill). (Mader S. *Inquiry into Life*, 6th edition, 1991; Chapter 7, p142).

Water availability and saturation levels in terrestrial environments vary considerably. Plant-associated bacteria experience different levels of hydration depending on the colonization site, prevailing climate conditions and soil composition. The phyllosphere and phylloplane (leaves and leaf surfaces) are relatively dry but can be wetted by rainfall and dew. The rhizosphere and rhizoplane (root microenvironment and root surfaces) and soil-borne seeds are more consistently hydrated with a surface water film that is highly dependent on soil saturation. Bacteria that can invade the internal plant vasculature experience the most consistent levels of saturation. Water limitation has dramatic effects on biofilm structure and, therefore, the saturation level of a particular environment and a specific tissue will profoundly affect biofilm growth [9–11*].

Among each major tissue type are a variety of microenvironments. For example, the surface characteristics vary along the length of the root [12]. Actively growing root tissues typically exhibit higher exudation rates into the soil, and root cap cells at the growing tip can be sloughed

away (Figure 1). Biofilms can be dramatically influenced by nutrient release and exudation at different sites. Leaf tissues often have a waxy cuticle that differs between the upper and lower portions of the leaf, interspersed with veins and petioles, trichomes and stomata. Phloem and xylem vessels are distinct tissue types within the vasculature that differ in fluid composition, architecture and spatial arrangement within leaves, stems and roots (Figure 1). Bacteria have adapted to each of these microenvironments, and the biofilms thus formed reflect the nature of their colonization sites.

Active and passive deposition on plant surfaces

Passive mechanisms of microbial deposition are common throughout the terrestrial environment, including wind and rain splash in the phyllosphere, and water flow in the rhizosphere [13]. Chemotaxis and motility are active mechanisms for establishment of biofilm communities [14–16]. Motility of several different pseudomonads appears to be important for competitive root colonization and long-term survival in soils [17–19]. Seed-associated

bacteria may also colonize the developing rhizosphere via chemotaxis and motility [17,18]. On leaf surfaces, passive deposition is important, as bacteria are removed and aerially transported to new locations [13]. Once deposited in the phyllosphere, motility may aid the bacteria in accessing a specific niche, but more often they congregate at their landing site [20]. In moist and nutrient-rich sites the bacteria can grow into aggregates and biofilms [20].


Biofilms in the rhizosphere

Root-associated pseudomonads have been studied extensively, and many of these promote the growth of host plants or are used as biocontrol agents [21]. Species of *Pseudomonas* form dense biofilms on both abiotic and biotic surfaces, and are a primary model in biofilm research [2]. *Pseudomonas putida* can respond rapidly to the presence of root exudates in soils, converging at root colonization sites and establishing stable biofilms [22[•]]. The plant-growth-promoting pseudomonads have been reported to discontinuously colonize the root surface, developing as small biofilms along epidermal fissures [23]. By contrast, recent studies analyzing pathogenic pseudomonads revealed dense, confluent biofilms on root surfaces [24[•],25^{••}]. Although the underlying cause for these different observations is unclear, it seems that pseudomonad root biofilms can range from relatively small multicellular clusters to extensive biofilm networks.

Azospirillum brasilense and related species are motile, heterotrophic proteobacteria that interact with roots of a variety of cereals such as wheat and maize, and often promote the growth of their host plants [26]. Although *A. brasilense* is a free-living nitrogen fixer, its ability to promote plant growth seems to be related to stimulation of root proliferation, rather than providing fixed nitrogen to the plant. The bacteria colonize root elongation zones and root hairs, forming dense biofilms [27]. Species of *Agrobacterium* and genera of symbiotic rhizobia not only cause neoplasia and symbiotic nodules on roots but are also effective root colonizers. Rhizobia preferentially associate with legume root hairs, stimulate root hair curling, infection thread elongation, and nodule formation on the appropriate host plant [28,29]. Microscopy of rhizobial cells within curled root hairs reveals small biofilm-type aggregates that provide the inocula for root invasion; the rhizobial cells migrate down infection threads as biofilm-like filaments towards the root interior (Figure 2a) [30[•]]. *Agrobacterium tumefaciens* and rhizobia can form dense, structurally complex biofilms on root surfaces, extensively coating the epidermis and root hairs, and these bacteria also form elaborate biofilms on abiotic surfaces (Figure 2b; AM Hirsch, personal communication) [31[•],32].

Gram-positive microbes also effectively colonize the rhizosphere and are well represented in soil populations [33]. Biocontrol agents such as *Bacillus cereus* develop dense surface-associated populations, and one recent study has

Figure 2



Colonization of plants by the *Rhizobiaceae*. (a) Curled root hair of alfalfa with red (DsRed) and green (GFP)-expressing *Sinorhizobium meliloti* in a mixed microcolony occupying the interior bend of the curl. The DsRed-labeled cells have initiated an infection thread. (Image courtesy of DJ Gage, [30[•]]). (b) Epifluorescence micrograph (Nikon E80040 X objective) of *Agrobacterium tumefaciens* C58 (P_{tac}-gfp) adhered to *Arabidopsis thaliana* Landsberg root segment. Overlay of *gfp* fluorescence and autofluorescence of plant tissue (T Danhorn and C Fuqua, unpublished).

linked biocontrol with the ability to form biofilms [25**]. Several functions known to influence biocontrol activity are also likely to play a role in biofilm formation [34].

A variety of specific functions are relevant to colonization and biofilm formation on plant roots. Motility via flagella or type IV pili is required for competitive colonization of roots by pseudomonads [16,19,35]. Motility mutants typically demonstrate only modest deficiencies in non-competitive root colonization, emphasizing the efficacy of passive rhizodeposition. Surface structures such as lipopolysaccharide and outer membrane proteins are also important in biofilm formation on roots. Hinsa *et al.* [36*] recently identified a 900-kDa cell surface protein called LapA (large adhesion protein A), that affects *P. fluorescens* colonization of glass, plastic and quartz sand, and is speculated to be a general adhesin. The LapA homologue in *P. putida* KT2440 is also required for competitive root colonization and seed adhesion [37]. LapA has domains that resemble adhesins involved in biofilm formation of Gram-positive bacteria, and a domain similar to Ca²⁺-binding proteins and haemolysins, often involved in host-cell interactions [36*]. For *Rhizobium leguminosarum* biovar *trifolii* a set of secreted agglutinins also thought to bind Ca²⁺ called Rap (*Rhizobium*-adhering) proteins localize to cell poles and are hypothesized to play a role in binding of rhizobial cells to plant tissues [38].

Production of exopolysaccharide is generally important in biofilm formation, and likewise can effect the interaction of bacteria with roots and root appendages [39,40]. Recent findings suggest that multiple polysaccharides modulate the chemical and physical attributes of the *P. aeruginosa* biofilm matrix on abiotic surfaces [41]. Such complexity may explain variable observations regarding the requirement for specific exopolysaccharides in biofilm formation and root association. For example, cellulose production in *A. tumefaciens* facilitates normal root adherence, and cellulose overproduction results in extremely dense biofilms ([42]; AG Matthyse, personal communication). By contrast, *A. tumefaciens* mutants that cannot synthesize the abundant exopolysaccharide succinoglycan (SCG) interact normally with roots, while mutants that overproduce SCG exhibit severely diminished adhesion (BE Ramey *et al.* unpublished data).

Later stages of biofilm maturation can also influence the structure of bacterial populations on roots. Recent findings suggest that *A. tumefaciens* biofilm formation on abiotic surfaces is regulated by the SinR transcription factor [31*]. SinR is a member of the FNR (fumarate and nitrate reductase) family of proteins, oxygen-responsive regulators that often control the transition to oxygen-limited conditions. Although it is unlikely that SinR senses oxygen directly, its expression is activated under oxygen limitation. *A. tumefaciens* with a *sinR* disruption formed a sparse, heterogeneous biofilm on abiotic sur-

faces, whereas strong *sinR* expression resulted in much thicker and less structured biofilms. The *sinR* biofilm phenotypes were recapitulated when examined on plant roots. Surface boundary layers and biofilms are generally oxygen-limited, and we speculate that SinR functions to respond to this feature of surface-associated growth. Other bacteria also appear to experience oxygen limitation during plant association [43,44].

Biofilms on seeds and sprouts

Bacterial adherence to seeds is a process that strongly influences rhizosphere colonization. Suppliers often deliberately coat their seed stocks with microbial biofilms to inoculate the developing rhizosphere. Additionally, biofilms on seeds and sprouts used for human consumption are common sources of infection. *P. putida* adheres effectively to seeds and will subsequently colonize the rhizosphere [37]. Several *P. putida* mutants, including one in the *lapA* homologue of *P. fluorescens*, are deficient in seed adherence and biofilm formation on inert surfaces, emphasizing the overlap between these activities. Recently, Coombs and Franco [45] identified endophytic populations of nonpathogenic actinobacteria in wheat tissues and determined that these were derived from interior colonization of surface-sterilized seeds. Endophytic seed populations help ensure future rhizosphere colonization.

Other studies of seed colonization have observed rod-shaped and coccid bacteria embedded within EPS in scanning electron micrographs of alfalfa seeds and sprouts [46,47]. Biofilms are notoriously resistant to washing and other common antibacterial treatments. Fett *et al.* found that both *Escherichia coli* O157:H7 and *Salmonella* populations on alfalfa sprouts required treatments much harsher than simple water washing to reduce the numbers of adherent microbes, and full removal was never achieved [48,49]. It seems likely that the surviving bacteria resided within biofilms, although this was not addressed.

Biofilm formation by vascular pathogens

Vascular pathogens inhabit the xylem or phloem of plant hosts and generally depend on insect vectors or wounding for dissemination. Several xylem-localized pathogens have received significant attention, while investigations of biofilm formation by phloem-restricted pathogens have focused primarily on spiroplasma in insects [50].

Xylella fastidiosa (*Xf*) is deposited into the xylem of plants by sap-feeding leafhoppers, and can induce Pierce's disease of grapevine and citrus variegated chlorosis [51]. Biofilms of *Xf* in the insect are composed of cells that are polarly attached to insect foregut tissue [52,53,54**]. In the plant host, most xylem vessels are sparsely colonized and asymptomatic, whereas densely populated vessels with biofilms are more rare, but symptomatic [54**]. Colonization depends on the *rpf* quorum-sensing (QS) system and the diffusible signal factor (DSF) [55*]. DSF


appears to govern expression of insect-specific adhesion factors [55[•]]. Insects exposed to a mutant in *rpfF*, encoding the DSF synthase, are not colonized by the bacteria and remain deficient for disease transmission. Interestingly, the *rpfF* mutant is hypervirulent when manually introduced into the plant host. Wild type *Xf* produces pit-membrane-degrading enzymes thought to aid movement into neighboring xylem vessels. It is plausible that expression of these enzymes is DSF-dependent. If so, a DSF mutant might densely populate xylem vessels, causing enhanced symptoms because it is unable to traverse the pit membrane [55[•]]. Fastidium exopolysaccharide is an important virulence factor for *Xf*. Bacteria attach to xylem vessels in the absence of the exopolysaccharide, while matrix-encased bacteria appear mainly in densely colonized vessels. Leite and co-workers proposed an adhesion model in which cell surface-exposed thiol groups associated with membrane features, impart a net negative cell surface charge, promoting divalent ion bridging for bacteria-to-bacteria and bacteria-to-host cell adhesion [56[•]]. The same group found that artificial media based on xylem chemistry stimulates *Xf* biofilm formation [57].

Xanthomonas campestris pv. *campestris* (*Xcc*) causes black rot on cruciferous plants, accessing the vasculature through wound sites in roots. Virulence involves degradative exoenzymes and the exopolysaccharide xanthan gum, both governed by *rpf*-encoded regulatory proteins and a DSF signal synthase [58[•]]. *Xcc* DSF was recently characterized as *cis*-11-methyl 2-dodecenoic acid, a novel α,β -unsaturated fatty acid QS signal [59]. DSF-dependent exopolysaccharide synthesis is necessary for biofilm formation and virulence, but not for bacterial adhesion. Candidate adhesion factors include pili and non-fimbrial adhesins. Dow *et al.* [58[•]] report that *Xcc* extracellular enzyme preparations induce dispersion of bacterial aggregates. The dispersion factor is an endo- β -(1,4)-mannanase (ManA) that expresses in an *rpf*/DSF-dependent manner and appears to facilitate spread of the pathogen through the plant vasculature.

Pantoea stewartii subsp. *stewartii* causes Stewart's wilt disease in maize and is transmitted by the corn flea beetle [60]. The bacteria reside primarily in the host xylem and produce large amounts of exopolysaccharide, controlled as a function of cell density through the EsaI/EsaR QS regulatory system [61,62]. Mutants of *esaI* and/or *esaR* alter bacterial adhesion, swarming motility, and biofilm formation [63]. Seedling infection assays using green fluorescent protein (GFP)-tagged strains show that the wild type strain colonizes the xylem vessels in discontinuous biofilms, while electron microscopic imaging shows bacterial aggregates covered with fibrous material associated with the xylem walls (Figure 3a,b).

Ralstonia solanacearum is a soil-borne pathogen that causes lethal wilt on many plants. Virulence depends on EPS and

Figure 3



Colonization of the vasculature by *Pantoea stewartii* subsp. *stewartii*. (a) GFP-tagged wild-type strain DC283 colonizing leaf xylem vessels of a susceptible maize cultivar. The discontinuous colonization pattern might be indicative of successive cycles of biofilm formation and dispersal as a strategy for systemic infection. Obtained using an Olympus IX70 inverted epifluorescence microscope, 40X magnification. (b) Scanning electron micrographs depicting colonization of sweetcorn xylem vessels. Image obtained on a LEO/Zeiss DSM 982 digital field emission scanning electron microscope (M Koutsoudis and SB von Bodman).

cell-wall-degrading enzymes controlled by a complex regulatory network [64[•]]. Denny and co-workers showed that the bacterium uses type IV pili for surface adhesion and twitching motility [64[•]]. Polar adhesion to plant cells is mediated by the PilA protein. A *pilA* mutant is less virulent and fails to form three-dimensional aggregates [64[•]]. Allen and colleagues [19] demonstrated a link between swimming motility and *Ralstonia* virulence and reported the observation of structures consistent with xylem biofilms.

Clavibacter michiganensis subsp. *sepedonicus* is a Gram-positive phytopathogen that causes bacterial ring rot in potato. Marques and colleagues [65] showed large bacterial, matrix-encased aggregates attached to the xylem vessels. Nonpathogenic Gram-positive filamentous actinobacteria were recently reported as endophytes of wheat [66[•]]. The bacteria were observed to form aggregates and

microcolonies in the intercellular spaces of healthy plant tissues, although it is not yet clear if they are disseminated through the vasculature.

Biofilms in epiphytic plant colonization

Aerial plant surfaces (i.e. the phylloplane) support large populations of bacterial epiphytes, including plant pathogens that multiply on the leaf surface before initiating disease [7[•],67]. The leaf surface is partitioned into preferred microhabitats along veins, near trichomes and stomates.

Pseudomonas syringae pv. *syringae* (*Pss*), the cause of brown spot disease on bean, colonizes the leaf surface sparsely in solitary small groups (fewer than ten cells), while larger populations (more than 1000 cells) primarily develop near trichomes or veins with higher nutrient availability. Large aggregates survive desiccation stress better than solitary cells [68]. Lindow and co-workers [11] report that the epiphytic fitness of *Pss* is governed by the AhII/AhIR QS system and the AefR regulator. AHL-null mutants are less tolerant to desiccations on leaves. Morris and colleagues assessed the epiphytic population structure of field-grown endive and cantaloupe and found that fluorescent pseudomonads were equally distributed in solitary or biofilm-associated populations, while Gram-positive epiphytes on cantaloupe tended to be in biofilms [6]. Leaf colonization by fluorescent pseudomonads may involve the deposition at new sites by solitary bacteria [48]. Large and small aggregates may vary as a function of nutrient availability at a given site [10].

Erwinia chrysanthemi (*Ech*) causes soft-rot disease through rapid maceration of plant tissue. Collmer and colleagues [69[•]] reported that *Ech* mutants in the filamentous hemagglutinin HecA have reduced leaf surface attachment and aggregate formation, fail to express macerating enzymes and are avirulent. The production of pectic enzymes may be QS-regulated, and therefore the inability to form bacterial aggregates may preclude pectinolytic enzyme secretion, illustrating how interference with an early phase of infection can dramatically impact successive steps.

Conclusions

Bacteria physically interact with plants in diverse ways. A common feature of this interaction is surface colonization, in which the microbes adhere to external and internal plant tissues as individual cells and in clusters. The adherent populations we define as biofilms exhibit a range of dimensions, locations and compositions. Each microenvironment of the plant has characteristic saturation levels, nutrient availabilities and surface chemistries, all of which strongly influence the form and activity of biofilms.

We have reviewed current examples of plant-associated biofilms, and some of the bacterial functions influencing

the establishment of these structures. A fundamental question in this regard is whether the process of biofilm formation *per se* drives or significantly impacts the dynamics of plant-microbe interactions and the effect of pathogens, symbionts and commensals on their hosts. *A priori*, the answer appears to be 'yes'. The number, conformation and viability of the associated bacteria must be important. A handful of recent studies such as those on *A. tumefaciens* and *X. fastidiosa* provide direct evidence to support this conjecture, but a great deal of research in different systems remains to determine how biofilm formation mechanisms are integrated with productive plant association.

Acknowledgements

We wish to thank our many colleagues who study plant-associated biofilms and apologize for those studies we were unable to discuss in this necessarily brief review. We also wish to thank AM Hirsch and DJ Gage for sharing unpublished results and biofilm images. Biofilm research is supported by the United States Department of Agriculture in the Fuqua laboratory (NRI 2002-35319-12636) and in the von Bodman laboratory (NRI 2002-35319-1237, and USDA Agricultural Experiment Station grant CNS00712).

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Webb JS, Givskov M, Kjelleberg S: **Bacterial biofilms: prokaryotic adventures in multicellularity.** *Curr Opin Microbiol* 2003, **6**:578-585.
 2. Parsek MR, Fuqua C: **Biofilms 2003: Emerging themes and challenges in studies of surface-associated microbial life.** *J Bacteriol* 2004, **186**:4427-4440.
 3. Stoodley P, Sauer K, Davies DG, Costerton JW: **Biofilms as complex differentiated communities.** *Annu Rev Microbiol* 2002, **56**:187-209.
 4. Sauer K, Camper AK, Ehrlich GD, Costerton JW, Davies DG: ***Pseudomonas aeruginosa* displays multiple phenotypes during development as a biofilm.** *J Bacteriol* 2002, **184**:1140-1154.
 5. Whiteley M, Bangera MG, Bumgarner RE, Parsek MR, Teitzel GM, Lory S, Greenberg EP: **Gene expression in *Pseudomonas aeruginosa* biofilms.** *Nature* 2001, **413**:860-864.
 6. Morris CE, Monier JM: **The ecological significance of biofilm formation by plant-associated bacteria.** *Annu Rev Phytopathol* 2003, **41**:429-453.
 7. Monier JM, Lindow SE: **Frequency, size, and localization of bacterial aggregates on bean leaf surfaces.** *Appl Environ Microbiol* 2004, **70**:346-355.
- This is a study of the spatial distribution of bacterial aggregates on leaf surfaces showing a non-random colonization pattern, with most bacteria associated in larger aggregates occupying preferred sites along potentially nutrient-rich leaf veins and glandular trichomes.
8. Bloemberg GV, Lugtenberg BJJ: **Bacterial biofilms on plants: relevance and phenotypic aspects.** In *Microbial Biofilms*. Edited by Ghannoum M, O'Toole GA. ASM Press; 2004.
 9. Chang WS, Halverson LJ: **Reduced water availability influences the dynamics, development, and ultrastructural properties of *Pseudomonas putida* biofilms.** *J Bacteriol* 2003, **185**:6199-6204.
 10. van de Mortel M, Halverson LJ: **Cell envelope components contributing to biofilm growth and survival of *Pseudomonas***

- putida** in low-water-content habitats. *Mol Microbiol* 2004, **52**:735-750.
11. Quinones B, Pujol J, Lindow SE: **Regulation of AHL production and its contribution to epiphytic fitness in *Pseudomonas syringae***. *Mol Plant Microbe Interact* 2004, **17**:521-531.
This is the first report that epiphytic fitness requires a functional quorum-sensing regulatory system.
 12. Walker TS, Bais HP, Grotewold E, Vivanco JM: **Root exudation and rhizosphere biology**. *Plant Physiol* 2003, **132**:44-51.
 13. Pedgley DE: **Aerobiology: The atmosphere as a source and sink for microbes**. In *Microbial Ecology of Leaves*. Edited by Andrews JH, Hirano SS. Springer-Verlag; 1991.
 14. Chesnokova O, Coutinho JB, Khan IH, Mikhail MS, Kado CI: **Characterization of flagella genes of *Agrobacterium tumefaciens*, and the effect of a bald strain on virulence**. *Mol Microbiol* 1997, **23**:579-590.
 15. O'Toole GA, Kolter R: **Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development**. *Mol Microbiol* 1998, **30**:295-304.
 16. Vande Broek A, Venderleyden J: **The role of bacterial motility, chemotaxis, and attachment in bacterial-plant interactions**. *Mol Plant Microbe Interact* 1995, **8**:800-810.
 17. Turnbull GA, Morgan JAW, Whipps JM, Saunders JR: **The role of bacterial motility in the survival and spread of *Pseudomonas fluorescens* in soil and in the attachment and colonisation of wheat roots**. *FEMS Microbiol Ecol* 2001, **36**:21-31.
 18. Turnbull GA, Morgan JAW, Whipps JM, Saunders JR: **The role of motility in the *in vitro* attachment of *Pseudomonas putida* PaW8 to wheat roots**. *FEMS Microbiol Ecol* 2001, **35**:57-65.
 19. Tans-Kersten J, Huang H, Allen C: ***Ralstonia solanacearum* needs motility for invasive virulence on tomato**. *J Bacteriol* 2001, **183**:3597-3605.
 20. Lindow SE, Leveau JH: **Phyllosphere microbiology**. *Curr Opin Biotechnol* 2002, **13**:238-243.
 21. Lugtenberg BJJ, Dekkers L, Bloemberg GV: **Molecular determinants of rhizosphere colonization by *Pseudomonas***. *Annu Rev Phytopathol* 2001, **39**:461-490.
 22. Espinosa-Urgel M, Kolter R, Ramos JL: **Root colonization by *Pseudomonas putida*: love at first sight**. *Microbiology* 2002, **148**:341-343.
This succinct and insightful study presents a method for real-time observation of corn root colonization by *Pseudomonas putida*. The study suggests rapid recognition and chemotaxis towards newly introduced root tissue followed by establishment and maintenance of biofilms.
 23. Bloemberg GV, Wijfjes AHM, Lamers GEM, Stuurman N, Lugtenberg BJJ: **Simultaneous imaging of *Pseudomonas fluorescens* WCS365 populations expressing three different autofluorescent proteins in the rhizosphere: new perspectives for studying microbial communities**. *Mol Plant Microbe Interact* 2000, **13**:1170-1176.
 24. Walker TS, Bais HP, Deziel E, Schweitzer HP, Rahme LG, Fall R, Vivanco JM: ***Pseudomonas aeruginosa*-plant root interactions. Pathogenicity, biofilm formations, and root exudation**. *Plant Physiol* 2004, **134**:3210-3331.
Sweet basil plants produce rosmarinic acid, an antibacterial substance upon infection by *Pseudomonas aeruginosa*, that protects pre-treated plants from infection, but cannot protect plants once the bacterial pathogen has had the opportunity to establish a biofilm. This study intriguingly reveals that biofilm formation can indeed protect invading pathogens against a chemical defense response from a plant host.
 25. Bais HP, Fall R, Vivanco JM: **Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production**. *Plant Physiol* 2004, **134**:307-319.
This work describes sensitive *in vitro* and in-soil plant infection models to study the biocontrol abilities of *B. subtilis* against *P. syringae* infection. The surfactin produced by *B. subtilis* is required for biofilm formation by that organism and is an antimicrobial agent. A surfactin-deficient mutant is ineffective in both biofilm formation and biocontrol against *P. syringae* infection of treated tomato plants.
 26. Burdman S, Okon Y, Jurkevitch E: **Surface characteristics of *Azospirillum brasilense* in relation to cell aggregation and attachment to plant roots**. *Crit Rev Microbiol* 2000, **26**:91-110.
 27. Assmus B, Hutzler P, Kirchhof G, Amann R, Lawrence JR, Hartmann A: ***In situ* localization of *Azospirillum brasilense* in the rhizosphere of wheat with fluorescently labeled rRNA-targeted oligonucleotide probes and scanning confocal laser microscopy**. *Appl Environ Microbiol* 1995, **61**:1013-1019.
 28. Smit G, Swart S, Lugtenberg BJJ, Kijne JW: **Molecular mechanisms of attachment of *Rhizobium* bacteria to plant roots**. *Mol Microbiol* 1992, **6**:2897-2903.
 29. Gage DJ: **Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia during nodulation of temperate legumes**. *Microb Mol Biol Rev* 2004, **68**:280-300.
 30. Gage DJ: **Analysis of infection thread development using Gfp- and DsRed-expressing *Sinorhizobium meliloti***. *J Bacteriol* 2002, **184**:7042-7046.
This beautiful confocal microscopy study of differentially-tagged *S. meliloti* follows the bacteria through colonization of alfalfa and infection thread formation. Dual infections yielded few mixed-population nodules, providing evidence that infection thread development is dependent on cell proliferation.
 31. Ramey BE, Matthyse AG, Fuqua C: **The FNR-type transcriptional regulator SinR controls maturation of *Agrobacterium tumefaciens* biofilms**. *Mol Microbiol* 2004, **52**:1495-1511.
This study identifies the SinR regulator of *Agrobacterium tumefaciens* that controls biofilm maturation on abiotic surfaces and root tissues. SinR is a component of a putative oxygen-limitation response pathway that may facilitate adaptation to the low oxygen levels within biofilms.
 32. Matthyse AG, McMahan S: **The effect of the *Agrobacterium tumefaciens attR* mutation on attachment and root colonization differs between legumes and other dicots**. *Appl Environ Microbiol* 2001, **67**:1070-1075.
 33. Boureau T, Jaques MA, Berruyer R, Dessaux Y, Dominguez H, Morris CE: **Comparison of the phenotypes and genotypes of biofilms and solitary epiphytic bacterial populations on broad-leaved endive**. *Microb Ecol* 2004, **47**:87-95.
 34. Dunn AK, Klimowicz AK, Handelsman J: **Use of a promoter trap to identify *Bacillus cereus* genes regulated by tomato seed exudate and a rhizosphere resident, *Pseudomonas aureofaciens***. *Appl Environ Microbiol* 2003, **69**:1197-1205.
 35. Lugtenberg BJJ, Chin AWTF, Bloemberg GV: **Microbe-plant interactions: principles and mechanisms**. *Antonie Van Leeuwenhoek* 2002, **81**:373-383.
 36. Hinsla SM, Espinosa-Urgel M, Ramos JL, O'Toole GA: **Transition from reversible to irreversible attachment during biofilm formation by *Pseudomonas fluorescens* WCS365 requires an ABC transporter and a large secreted protein**. *Mol Microbiol* 2003, **49**:905-918.
This paper describes genetic and biochemical analysis of LapA, a fascinating multidomain, 900-kDa protein from *P. fluorescens* WCS365, which localized to the cell surface and was required for irreversible attachment to abiotic surfaces. LapA is similar to a *P. putida* protein shown to be required for seed attachment, root colonization and biofilm formation.
 37. Espinosa-Urgel M, Salido A, Ramos JL: **Genetic analysis of functions involved in adhesion of *Pseudomonas putida* to seeds**. *J Bacteriol* 2000, **182**:2363-2369.
 38. Ausmees N, Jacobsson K, Lindberg M: **A unipolarly located, cell-surface-associated agglutinin, RapA, belongs to a family of *Rhizobium*-adhering proteins (Rap) in *Rhizobium leguminosarum* bv. *trifolii***. *Microbiol SGM* 2001, **147**:549-559.
 39. Bianciotto V, Andreotti S, Balestrini R, Bonfante P, Perotto S: **Extracellular polysaccharides are involved in the attachment of *Azospirillum brasilense* and *Rhizobium leguminosarum* to arbuscular mycorrhizal structures**. *Eur J Histochem* 2001, **45**:39-49.
 40. Bianciotto V, Andreotti S, Balestrini R, Bonfante P, Perotto S: **Mucoid mutants of the biocontrol strain *Pseudomonas fluorescens* CHA0 show increased ability in biofilm formation**

- on mycorrhizal and non-mycorrhizal carrot roots. *Mol Plant Microbe Interact* 2001, **14**:255-260.
41. Friedman L, Kolter R: **Genes involved in matrix formation in *Pseudomonas aeruginosa* PA14 biofilms.** *Mol Microbiol* 2004, **51**:675-690.
 42. Matthyse AG: **Role of bacterial cellulose fibrils in *Agrobacterium tumefaciens* infection.** *J Bacteriol* 1983, **154**:906-915.
 43. Zhulin IG, Bespalov VA, Johnson MS, Taylor BL: **Oxygen taxis and proton motive force in *Azospirillum brasilense*.** *J Bacteriol* 1996, **178**:5199-5204.
 44. Okinaka Y, Yang C-H, Perna NT, Keen NT: **Microarray profiling of *Erwinia chrysanthemi* 3937 genes that are regulated during plant infection.** *Mol Plant Microbe Interact* 2002, **15**:619-629.
 45. Coombs JT, Franco CMM: **Isolation and identification of actinobacteria from surface-sterilized wheat roots.** *Appl Environ Microbiol* 2003, **69**:5603-5608.
 46. Fett WF, Cooke PH: **Scanning electron microscopy of native biofilms on mung bean sprouts.** *Can J Microbiol* 2003, **49**:45-50.
 47. Matos A, Garland JL, Fett WF: **Composition and physiological profiling of sprout-associated microbial communities.** *J Food Prot* 2002, **65**:1903-1908.
 48. Fett WF, Cooke PH: **Reduction of *Escherichia coli* O157: H7 and *Salmonella* on laboratory-inoculated alfalfa seed with commercial citrus-related products.** *J Food Prot* 2003, **66**:1158-1165.
 49. Fett WF: **Reduction of *Escherichia coli* O157: H7 and *Salmonella* spp. on laboratory-inoculated Mung bean seed by chlorine treatment.** *J Food Prot* 2002, **65**:848-852.
 50. Ammar E-D, Fulon D, Bai X, Meulia T, Hogenhout SA: **An attachment tip and pili-like structures in insect- and plant-pathogenic spiroplasmas of the class of Mollicutes.** *Arch Microbiol* 2004, **181**:97-105.
 51. Purcell AH, Hopkins DL: **Fastidious xylem-limited bacterial pathogens.** *Annu Rev Phytopathol* 1996, **34**:131-151.
 52. da Silva FR, Vettore AL, Kemper EL, Leite B, Arruda P: **Fastidium gum: the *Xylella fastidiosa* exopolysaccharide involved in bacterial pathogenicity.** *FEMS Microbiol Lett* 2001, **203**:165-171.
 53. Marques LLR, Ceri H, Manfio GP, Reid DM, Olsen ME: **Characterization of biofilm formation by *Xylella fastidiosa* in vitro.** *Plant Dis* 2002, **86**:633-638.
 54. Newman KL, Almeida RP, Purcell AH, Lindow SE: **Use of a green fluorescent strain for analysis of *Xylella fastidiosa* colonization of *Vitis vinifera*.** *Appl Environ Microbiol* 2003, **69**:7319-7327.
- Confocal laser-scanning microscopy of GFP-tagged *Xylella fastidiosa* cells shows that the bacteria generally colonize xylem vessels in small aggregates and that rare blockages can be linked to symptom development. More importantly, this study shows that movement of bacteria through bordered pits into adjacent vessels is a key strategy of colonization.
55. Newman KL, Almeida RP, Purcell AH, Lindow SE: **Cell-cell signaling controls *Xylella fastidiosa* interactions with both insects and plants.** *Proc Natl Acad Sci USA* 2004, **101**:1737-1742.
- This timely report indicates that insect vector colonization and transmission of *Xylella fastidiosa* is dependent on intercellular communication through the diffusible signal factor, and that this signaling may limit virulence in the infected plant.
56. Leite B, Ishida ML, Alves E, Carrier H, Pascholati SF, Kitajima EW: **Genomic and X-ray microanalysis indicate that Ca²⁺ and thiols mediate the aggregation and adhesion of *Xylella fastidiosa*.** *Braz J Med Biol Res* 2002, **35**:645-650.
- These authors offer a model for bacteria-bacteria and bacteria-plant-cell interactions based on divalent cation bridging between negatively charged bacterial-surface thiols and nonspecific negatively charged moieties of the xylem wall.
57. Leite B, Anderson PC, Ishida ML: **Colony aggregation and biofilm formation in xylem chemistry-based media for *Xylella fastidiosa*.** *FEMS Microbiol Lett* 2004, **230**:283-290.
 58. Dow JM, Crossman L, Findlay K, He Y-Q, Feng J-X, Tang J-L: **Biofilm dispersal in *Xanthomonas campestris* is controlled by cell-cell signaling and is required for full virulence to plants.** *Proc Natl Acad Sci USA* 2003, **100**:10995-11000.
- In this intriguing study, cell-cell signaling via the diffusible signal factor is shown to stimulate dispersal of *Xanthomonas campestris* pv. *campestris* aggregates through the activity of an endo- β -(1,4)-mannanase enzyme, ManA. A *manA* mutant demonstrated attenuated virulence.
59. Wang LH, He Y, Gao Y, Wu JE, Dong YH, He C, Weng LX, Xu JL, Tay L, Fang RX *et al.*: **A bacterial cell-cell communication signal with cross-kingdom structural analogues.** *Mol Microbiol* 2004, **51**:903-912.
 60. Leigh JA, Coplin DL: **Exopolysaccharide in plant-bacterial interactions.** *Annu Rev Microbiol* 1992, **46**:307-346.
 61. von Bodman SB, Majerczak DR, Coplin DL: **A negative regulator mediates quorum-sensing control of exopolysaccharide production in *Pantoea stewartii* subsp. *stewartii*.** *Proc Natl Acad Sci USA* 1998, **95**:7687-7692.
 62. von Bodman SB, Bauer WD, Coplin DL: **Quorum sensing in plant-pathogenic bacteria.** *Annu Rev Phytopathol* 2003, **41**:455-482.
 63. von Bodman SB, Koutsoudis M, Minogue TD: **Steps in biofilm formation and infection in *Pantoea stewartii* subsp. *stewartii*.** *Phytopathology* 2003, **92**:S97.
 64. Kang Y, Liu H, Genin S, Schell MA, Denny TP: ***Ralstonia solanacearum* requires type 4 pili to adhere to multiple surfaces and for natural transformation and virulence.** *Mol Microbiol* 2002, **46**:427-437.
- This is a report of twitching motility, autoaggregation and biofilm formation for *Ralstonia solanacearum*. The authors present evidence for the role of type IV pili in plant pathogenesis.
65. Marques LLR, De Boer SH, Ceri H, Olsen ME: **Evaluation of biofilms formed by *Clavibacter michiganensis* subsp. *sepedonicus*.** *Phytopathology* 2003, **93**:S57.
 66. Coombs JT, Franco CMM: **Visualization of an endophytic *Streptomyces* species in wheat seed.** *Appl Environ Microbiol* 2003, **69**:4260-4262.
- This paper provides striking microscopic evidence for the endophytic association of a *Streptomyces*-type actinobacteria with wheat seeds and primordial tissues.
67. Beattie GA, Lindow SE: **Bacterial colonization of leaves: a spectrum of strategies.** *Appl Environ Microbiol* 1999, **60**:3799-3808.
 68. Monier JM, Lindow SE: **Differential survival of solitary and aggregated bacterial cells promotes aggregate formation on leaf surfaces.** *Proc Natl Acad Sci USA* 2003, **100**:15977-15982.
 69. Rojas CM, Ham JH, Deng WL, Doyle JJ, Collmer A: ***HecA*, a member of a class of adhesins produced by diverse pathogenic bacteria, contributes to the attachment, aggregation, epidermal cell killing, and virulence phenotypes of *Erwinia chrysanthemi* EC16 on *Nicotiana glauca* seedlings.** *Proc Natl Acad Sci USA* 2002, **99**:13142-13147.
- The *hecA* gene encodes a filamentous hemagglutinin-like adhesin, and is shown to be a broadly important virulence factor in both plant and animal pathogens. This work also demonstrates that interference with an early phase of infection can dramatically impact subsequent steps of infection.